


Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently amended) A compound having the formula (I):




(I)

or a stereoisomer or a pharmaceutically acceptable salt thereof, wherein:

A is C₁₋₆ alkylene; R, R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, S-alkyl or alkylthiol, X is >C₁₋₆ alkylene, >C=O or >C=S or a single bond; and Y is halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, ~~alkoxy~~, carboxy, carbonylamido, or

styryl, wherein where Y is a ring it may be ring-substituted by up to four substituents independently selected from among hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, S-alkyl, alkylthiol, or -COQ, where Q is hydroxy, C₁₋₆ alkoxy, amino, mono- C₁₋₆ alkylamino, di- C₁₋₆ alkylamino, hydroxylamino, C₁₋₄ alkoxyamino or aryl-C₁₋₄-alkoxyamino; but excluding (a) the compounds where simultaneously X is >C=O, Y is methyl, A is CH₂CH₂, R is 5-methoxy, R₁ is H or formyl and R₂ is H, (b) the compounds where the moiety -A(R₂)-NH-X-Y is -CH₂CH(COQ)-NH₂ or -CH(haloalkyl)-CH(COQ)-NH₂, and (c) the compounds where simultaneously X is a single bond, Y is arylalkyl, A is CH₂CH₂CH₂, both R₁ and R₂ are H and R is 4-halo where the moiety -CO-A(R₂)-NH-X-Y is deemed to be in the 1-position of the depicted benzene ring.

2. (Currently amended) A compound according to claim 1, having formula (II):



(II)

wherein R is hydrogen, methyl or methoxy, R₁ is hydrogen or formyl, R₂ is hydrogen or carboxyl, and R₃ is hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, S-alkyl or alkylthiol, or a ~~and~~ stereoisomers or ~~and~~ pharmaceutically acceptable salts thereof.

3. (Currently amended) A compound ~~Compounds~~ according to claim 1, where in formula (I), Y is 2-furyl, 2-dihydrofuryl, or 2-tetrahydrofuryl, any of which may be substituted by 1-2 substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, OH, nitro, or Y is styryl which is ring-substituted by up to two substituents independently selected from among halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, OH, nitro, aryl, aryl-C₁₋₄ alkyl, or aryl-C₁₋₄ alkoxy, and stereoisomers and pharmaceutically acceptable salts thereof.

4. (Currently amended) A compound ~~Compounds~~ according to claim 1, or a ~~and stereoisomers or and~~ pharmaceutically acceptable salts thereof, where in formula (I), R₂ is hydrogen and at least one of the following conditions applies, namely:

R is 5-methoxy; or

A is CH₂CH₂ ~~or CH₂CH~~COOH; or

R₁ is hydrogen; or

X is a single bond and Y is a 2,4-dinitrophenyl group.

5. (Currently amended) A compound ~~Compounds~~ according to claim 1, or a ~~and stereoisomers or and~~ pharmaceutically acceptable salts thereof, where in formula (I), X and Y are selected in combination as follows:

X is -CO- and Y is 2-furyl; or

X is -CO- and Y is 2-tetrahydrofuryl; or

X is -CH₂- and Y is 2-tetrahydrofuryl; or

X is -CO- and Y is 2-acetoxyphenyl; or

X is -CO- and Y is 3,4-dihydroxystyryl or 3,4-dihydroxycinnamoyloxy.

6. (Currently amended) A compound ~~Compounds~~ according to claim 5, wherein at least one of the following conditions applies, namely:

R is 5-methoxy; or

A is CH₂CH₂ or

A-R₂ is CH₂CHCOOH; or

R₁ is hydrogen.

7. (Original) A compound according to claim 1, which is 3-(2-aminobenzoyl)-2-(2,4-dinitroanilino)propanoic acid, and stereoisomers and pharmaceutically acceptable salts thereof.

8. (Previously presented) A compound according to claim 1, which is 2-(2-aminobenzoyl)-N-(2,4-dinitrophenyl)ethylamine, or a pharmaceutically acceptable salt thereof.

9. (Original) A pharmaceutical formulation containing a therapeutically effective amount of at least one compound as defined in claim 1 in association with at least one

pharmaceutically acceptable ingredient selected from diluents, preservatives, solubilizers, emulsifiers, adjuvants, excipients and carriers.

10. (Original) A pharmaceutical formulation according to claim 9, which is further characterized by at least one of the following features:

- (i) it is adapted for oral, rectal, parenteral, transbuccal, intrapulmonary or transdermal administration;
- (ii) it is in unit dosage form, each unit dosage comprising an amount of said at least one compound which lies within the range of 0.0025-1000 mg;
- (iii) it is a controlled release formulation, wherein said at least one compound is released at a predetermined controlled rate;
- (iv) it comprises additionally at least one known therapeutically active ingredient selected from neuroleptics, thymoleptics, anxiolytics, tranquilizers, analgesics, and antiparkinson's drugs.

11. (Previously presented) A method of treating a subject suffering from a physiological condition selected from the group consisting of stroke, ischemia, CNS trauma, hypoglycemia and surgery, CNS disorders, overstimulation of the excitatory amino acids, psychiatric disorders, epilepsy or other convulsive disorders, anxiety, psychosis, senile dementia, multi-infarct dementia, chronic pain (analgesia), glaucoma, CMV retinitis, urinary incontinence, impotence, cardiovascular disorders, blood

coagulation, neuropathy, anti-inflammatory, chronobiological-related disorders, seasonal-related disorders, endocrine indications, precocious puberty, premenstrual syndrome, hyperprolactinemia, growth hormone deficiency, neoplastic disease, benign or tumor prostate growth, immune system disorders, conditions associated with senescence, ophthalmological diseases, cluster headache, migraine, or weight gain disorders, which comprises administering a therapeutically effective amount of a compound of formula I or a stereoisomer or a pharmaceutically acceptable salt thereof as defined in claim 1.

12. (Currently amended) The method of claim 11, wherein said compound or stereoisomer or salt is administered in association with at least one pharmaceutically acceptable ingredient selected from diluents, preservatives, solubilizers, emulsifiers, adjuvants, excipients and carriers ~~in a pharmaceutical formulation as defined in claim 9.~~

13. (Currently amended) The method of claim 12, wherein said pharmaceutical formulation is further characterized by at least one of the following features:

(i) it is adapted for oral, rectal, parenteral, transbuccal, intrapulmonary or transdermal administration;


(ii) it is in unit dosage form, each unit dosage comprising an amount of said at least one compound which lies within the range of 0.0025-1000 mg;

(iii) it is a controlled release formulation, wherein said at least one compound is released at a predetermined controlled rate;

(iv) it comprises additionally at least one known therapeutically active ingredient selected from neuroleptics, thymoleptics, anxiolitics, tranquilizers, analgesics, and anti-Parkinson's drugs ~~is as defined in claim 10.~~

14. (Previously presented) A method for regulating fertility, puberty or pelage color in animal breeding, which comprises administering to a breeding animal an effective amount of a compound of formula I or a stereoisomer or pharmaceutically acceptable salt as defined in claim 1.

15. (Currently amended) A compound having the formula (I):




(I)

or a stereoisomer or a pharmaceutically acceptable salt thereof, wherein:

A is C₁₋₆ alkylene; R, R₁ and R₂ are independently hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, S-alkyl or alkylthiol, X is >C₁₋₆ alkylene, >C=O or >C=S; and Y is hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, or styryl, wherein where Y is a ring it may be ring-substituted by up to four substituents

independently selected from among hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, ~~alkoxy~~, S-alkyl, alkylthiol or -COQ, where Q is hydroxy, C₁₋₆ alkoxy, amino, mono- C₁₋₆ alkylamino, di- C₁₋₆ alkylamino, hydroxylamino, C₁₋₄ alkoxyamino or aryl-C₁₋₄-alkoxyamino; but excluding (a) the compounds where simultaneously X is >C=O, Y is methyl, A is CH₂CH₂, R is 5-methoxy, R₁ is H or formyl and R₂ is H and (b) the compounds where the moiety -A(R₂)-NH-X-Y is -CH₂CH(COQ)-NH₂ or -CH(haloalkyl)-CH(COQ)-NH₂.

16. (Previously presented) A compound having the formula (I):



(I)

or a stereoisomer or a pharmaceutically acceptable salt thereof, wherein:

A is C₁₋₆ alkylene; R, R₁ and R₂ are independently halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, S-alkyl or alkylthiol, and each of R₁ and R₂ independently also can be hydrogen, X is >C₁₋₆ alkylene, >C=O, >C=S or a single bond; and Y is hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido, or styryl, wherein where Y is a ring it may be ring-

substituted by up to four substituents independently selected from among hydrogen, halo, haloalkyl, aryl, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, hydroxyalkyl, nitro, amino, cyano, cyanamido, guanidine, amidino, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, S-alkyl, alkylthiol or -COQ, where Q is hydroxy, C₁₋₆ alkoxy, amino, mono- C₁₋₆ alkylamino, di- C₁₋₆ alkylamino, hydroxylamino, C₁₋₄ alkoxyamino or aryl-C₁₋₄-alkoxyamino; but excluding (a) the compounds where simultaneously X is >C=O, Y is methyl, A is CH₂CH₂, R is 5-methoxy, R₁ is H or formyl and R₂ is H, (b) the compounds where the moiety -A(R₂)-NH-X-Y is -CH₂CH(COQ)-NH₂ or -CH(haloalkyl)-CH(COQ)-NH₂, and (c) the compounds where simultaneously X is a single bond, Y is arylalkyl, A is CH₂CH₂CH₂, both R₁ and R₂ are H and R is 4-halo where the moiety -CO-A(R₂)-NH-X-Y is deemed to be in the 1-position of the depicted benzene ring.

17. (Currently amended) A compound according to claim 15 or 16, where in formula (I), Y is 2-furyl, 2-dihydrofuryl or 2-tetrahydrofuryl, any of which may be substituted by 1-2 substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, OH, nitro, or Y is hydrogen or styryl which is ring-substituted by up to two substituents independently selected from among halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, OH, nitro, aryl, aryl-C₁₋₄ alkyl, or aryl-C₁₋₄ alkoxy, or a ~~and~~ stereoisomers or ~~and~~ pharmaceutically acceptable salts thereof.

18. (New) A compound according to claim 1, or a stereoisomer or pharmaceutically acceptable salt thereof, where in formula (I) A-R₂ is CH₂CHCOOH.

19. (New) A pharmaceutical formulation containing a therapeutically effective amount of at least one compound as defined in claim 15 or 16 in association with at least one pharmaceutically acceptable ingredient selected from diluents, preservatives, solubilizers, emulsifiers, adjuvants, excipients and carriers.

20. (New) A pharmaceutical formulation according to claim 19, which is further characterized by at least one of the following features:

- (i) it is adapted for oral, rectal, parenteral, transbuccal, intrapulmonary or transdermal administration;
- (ii) it is in unit dosage form, each unit dosage comprising an amount of said at least one compound which lies within the range of 0.0025-1000 mg;
- (iii) it is a controlled release formulation, wherein said at least one compound is released at a predetermined controlled rate;
- (iv) it comprises additionally at least one known therapeutically active ingredient selected from neuroleptics, thymoleptics, anxiolytics, tranquilizers, analgesics, and antiparkinson's drugs.

21. (New) A method of treating a subject suffering from a physiological condition selected from the group consisting of stroke, ischemia, CNS trauma, hypoglycemia and surgery, CNS disorders, overstimulation of the excitatory amino acids, psychiatric disorders, epilepsy or other convulsive disorders, anxiety, psychosis, senile dementia, multi-infarct dementia, chronic pain (analgesia), glaucoma, CMV retinitis, urinary incontinence, impotence, cardiovascular disorders, blood coagulation, neuropathy, anti-inflammatory, chronobiological-related disorders, seasonal-related disorders, endocrine indications, precocious puberty, premenstrual syndrome, hyperprolactinemia, growth hormone deficiency, neoplastic disease, benign or tumor prostate growth, immune system disorders, conditions associated with senescence, ophthalmological diseases, cluster headache, migraine, or weight gain disorders, which comprises administering a therapeutically effective amount of a compound of formula I or a stereoisomer or a pharmaceutically acceptable salt thereof as defined in claim 15 or 16.

22. (New) The method of claim 21, wherein said compound or stereoisomer or salt is administered in association with at least one pharmaceutically acceptable ingredient selected from diluents, preservatives, solubilizers, emulsifiers, adjuvants, excipients and carriers.

23. (New) The method of claim 22, wherein said pharmaceutical formulation is further characterized by at least one of the following features:

- (i) it is adapted for oral, rectal, parenteral, transbuccal, intrapulmonary or transdermal administration;
- (ii) it is in unit dosage form, each unit dosage comprising an amount of said at least one compound which lies within the range of 0.0025-1000 mg;
- (iii) it is a controlled release formulation, wherein said at least one compound is released at a predetermined controlled rate;
- (iv) it comprises additionally at least one known therapeutically active ingredient selected from neuroleptics, thymoleptics, anxiolitics, tranquilizers, analgesics, and anti-Parkinson's drugs.

24. (New) A method for regulating fertility, puberty or pelage color in animal breeding, which comprises administering to a breeding animal an effective amount of a compound of formula I or a stereoisomer or pharmaceutically acceptable salt as defined in claim 15 or 16.